

AARC Clinical Practice Guideline

Nasotracheal Suctioning—2004 Revision & Update

NTS 1.0 PROCEDURE:

Nasotracheal suctioning (NTS) for tracheal aspiration is a component of resuscitation and bronchial hygiene therapy.¹

NTS 2.0 DESCRIPTION/DEFINITION:

NTS is intended to remove accumulated saliva, pulmonary secretions, blood, vomitus, and other foreign material from the trachea and nasopharyngeal area that cannot be removed by the patient's spontaneous cough or other less invasive procedures. NTS has been used to maintain a patent airway thus ensuring adequate oxygenation and ventilation² and avoiding intubation that was solely intended for the removal of secretions.^{1,3-6}

NTS refers to the insertion of a suction catheter through the nasal passage and pharynx into the trachea without a tracheal tube or tracheostomy (although a nasopharyngeal airway may be used)^{3,7} in order to aspirate accumulated secretions or foreign material.^{1,4,6}

The clearance of secretions is accomplished by application of subatmospheric pressure applied to a sterile, flexible, multi-eyed catheter^{1,6,8-10} on withdrawal only.⁹⁻¹⁵ Appropriate subatmospheric pressures are

- 2.1 Neonates: 60-80 mm Hg^{16,17}
- 2.2 Infants: 80-100 mm Hg^{16,18}
- 2.3 Children: 100-120 mm Hg^{13,16}
- 2.4 Adults: 100-150 mm Hg^{1,19}

Negative pressures should not exceed 150 mm Hg as higher pressures have been shown to cause trauma, hypoxemia and atelectasis.^{3,12,14,20-25}

NTS 3.0 SETTINGS:

NTS is performed in a wide variety of settings, and this guideline applies to patients of all ages.

- 3.1 Critical care¹
- 3.2 Emergency room or department¹

3.3 Inpatient acute care¹

3.4 Extended care and skilled nursing facility care¹

3.5 Home care¹

3.6 Outpatient or ambulatory care

NTS 4.0 INDICATIONS:

The need to maintain a patent airway and remove saliva, pulmonary secretions, blood, vomitus, or foreign material from the trachea in the presence of

4.1 Inability to clear secretions when audible or visible evidence of secretions in the large/central airways that persist in spite of patient's best cough effort.^{1,5,6,25-27} This is evidenced by one or more of the following

4.1.1 Visible secretions in the airway^{1,27}

4.1.2 Chest auscultation of coarse, gurgling breath sounds, rhonchi^{1,14,21,27,28} or diminished breath sounds²¹

4.1.3 Feeling of secretions in the chest (increased tactile fremitus)²¹

4.1.4 Suspected aspiration of gastric or upper airway secretions¹

4.1.5 Clinically apparent increased work of breathing¹

4.1.6 Deterioration of arterial blood gas values suggesting hypoxemia or hypercarbia^{1,21}

4.1.7 Chest radiographic evidence of retained secretions resulting in atelectasis or consolidation^{1,27}

4.1.8 Restlessness^{21,28}

4.2 To stimulate cough^{1,2,27,29} or for unrelieved coughing²¹

4.3 To obtain a sputum sample for microbiological or cytological analysis^{1,2,27}

NTS 5.0 CONTRAINDICATIONS:

Listed contraindications are relative unless marked as absolute.

5.1 Occluded nasal passages^{1,6}

- 5.2 Nasal bleeding¹
- 5.3 Epiglottitis or croup (absolute)^{1,6}
- 5.4 Acute head, facial, or neck injury^{1,2,6}
- 5.5 Coagulopathy or bleeding disorder^{1,3,6}
- 5.6 Laryngospasm^{1,3,6}
- 5.7 Irritable airway¹
- 5.8 Upper respiratory tract infection¹
- 5.9 Tracheal surgery⁶
- 5.10 Gastric surgery with high anastomosis⁶
- 5.11 Myocardial infarction⁶
- 5.12 Bronchospasm²

NTS 6.0 HAZARDS/COMPLICATIONS:

- 6.1 Mechanical trauma (mucosal hemorrhage, tracheitis, epistaxis from laceration of nasal turbinates, and perforation of the pharynx)^{1,6,14,17,26,27,30-34}
 - 6.1.1 Laceration of nasal turbinates^{8,35}
 - 6.1.2 Perforation of the pharynx³⁶
 - 6.1.3 Nasal irritation/bleeding⁷
 - 6.1.4 Tracheitis^{1,17}
 - 6.1.5 Mucosal hemorrhage^{2,32}
 - 6.1.6 Uvular edema³⁷
- 6.2 Hypoxia/hypoxemia^{1,2,6,17,27,33,38-41}
- 6.3 Cardiac dysrhythmias/arrest^{2,4,6,14,33-35}
- 6.4 Bradycardia^{1,2,6,27,38,41-44}
- 6.5 Increase in blood pressure^{1,2,6,38,40,45}
- 6.6 Hypotension^{1,38}
- 6.7 Respiratory arrest³⁵
- 6.8 Uncontrolled coughing^{1,2,7,34}
- 6.9 Gagging/vomiting^{1,6,7,46}
- 6.10 Laryngospasm^{1,3,35}
- 6.11 Bronchoconstriction/bronchospasm^{1,14,33,34}
- 6.12 Discomfort^{7,41} and pain^{1,2,7,41}
- 6.13 Nosocomial infection^{1,2,27,34,44}
- 6.14 Atelectasis^{2,8,14,17,27,33}
- 6.15 Misdirection of catheter^{6,7,34}
- 6.16 Increased intracranial pressure (ICP)^{6,28,40,41,45,47-49}
 - 6.16.1 Intraventricular hemorrhage^{14,40,50}
 - 6.16.2 Exacerbation of cerebral edema
- 6.17 Pneumothorax¹⁷

NTS 7.0 LIMITATIONS OF METHOD:

- 7.1 NTS is a blind procedure with inherent risks (refer to complications).^{1,6,7,44}
- 7.2 Risks are increased in a combative or uncooperative patient.
- 7.3 Duration of application of subatmospheric pressure, or suction, should be limited to no

greater than 15 seconds.^{1,2,6,9,13,14,17,20,21,24,28,39,51}

7.4 Controversy exists concerning possible overuse of this procedure.^{8,14,26,52}

NTS 8.0 ASSESSMENT OF NEED:

8.1 Personnel should perform a baseline assessment for indications of respiratory distress and the need for NTS as recognized by presenting indications as listed above. This should include but not be limited to

- 8.1.1 Auscultation of chest^{1,3,9,12,14,27,53,54}
- 8.1.2 Monitor patient's heart rate^{3,12,14}
- 8.1.3 Respiratory rate¹²
- 8.1.4 Cardiac rhythm^{12,14}
- 8.1.5 Oxygen saturation^{12,14}
- 8.1.6 Skin color and perfusion¹²
- 8.1.7 Personnel should assess effectiveness of cough¹

8.2 Prepare the patient for the procedure by providing an appropriate explanation along with adequate sedation and pain relief as needed.^{2,9,12}

NTS 9.0 ASSESSMENT OF OUTCOME:

Effectiveness of NTS should be reflected by assessing patient post suction for

- 9.1 Improved breath sounds^{1,36}
- 9.2 Removal of secretions^{1,36}
- 9.3 Improved blood gas data or pulse oximetry¹
- 9.4 Decreased work of breathing (decreased respiratory rate or dyspnea)¹

NTS 10.0 RESOURCES:

10.1 Equipment:

- 10.1.1 Vacuum source^{1,6,27}
- 10.1.2 Calibrated, adjustable regulator^{1,55}
- 10.1.3 Collection vessel and connecting tubing¹
- 10.1.4 Sterile, flexible, multiple-eyed suction catheter^{1,6,8-10} of appropriate caliber^{1,6,10,27,52,54}
- 10.1.5 Sterile disposable gloves^{1,6,27,52}
- 10.1.6 Sterile water and cup^{1,52}
- 10.1.7 Water-based lubricant^{1,6,27,52} and/or normal saline^{1,52}
- 10.1.8 Local anesthetic is sometimes used to reduce discomfort¹
- 10.1.9 Nasopharyngeal airway when frequent NTS is required^{1,3,7,54}
- 10.1.10 Resuscitation bag with mask^{1,6,27,35,56}

In the acute care setting, with initiation of NTS or when working with the unstable patient, the following are recommended

10.1.11 Electrocardiogram monitor^{1,27}

10.1.12 Oxygen (hyperoxygenation with appropriate delivery device as indicated)^{1,6,14,27,28,39,41,51,57,58}

10.1.13 Personnel protective equipment for Standard Precautions^{1,12,20,23,59-62}

10.1.14 Stethoscope²⁷

10.2 Personnel:

10.2.1 Level I caregiver may be the provider of service *after* Level II personnel have established need by patient assessment and the first NTS episode has been completed. Level I personnel must demonstrate

10.2.1.1 Knowledge of proper assembly and use of equipment²

10.2.1.2 Knowledge of upper airway anatomy and physiology^{35,44}

10.2.1.3 Ability to recognize secretion retention on auscultation^{1,3,12,27,54}

10.2.1.4 Ability to monitor vital signs and assess patient's condition and response to procedure^{3,12,14}

10.2.1.5 Ability to recognize and respond to adverse reactions and complications of procedures

10.2.1.6 Ability to employ technique of cardiopulmonary resuscitation when indicated

10.2.1.7 Ability to evaluate and document procedure effectiveness and patient response

10.2.2 Level II provider initially assesses the patient, determines the need for NTS, and evaluates response to and effectiveness of first episode. Level II personnel have all the skills of Level I providers plus:

10.2.2.1 Knowledge and understanding of patient's disease, goals, and limitation of NTS⁴⁴

10.2.2.2 Recognition and understanding of basis of pathophysiology

10.2.2.3 Ability to perform initial treatment and be available to troubleshoot the procedure

10.2.2.4 Ability to modify techniques and equipment and take definitive action in response to adverse reaction

10.2.2.5 Ability to detect adverse reactions and avoid patient harm by employing techniques of cardiopulmonary resuscitation with mechanical airway adjuncts and bag-mask devices

10.2.2.6 Knowledge of basic electrocardiogram and dysrhythmia recognition

10.2.2.7 Knowledge of signs and symptoms of decreased cardiac output, oxygenation, and perfusion

10.2.2.8 Ability to teach Level I and lay personnel providing home care

10.2.3 Home care should be provided by lay personnel trained and knowledgeable in

10.2.3.1 Proper assembly and use of equipment

10.2.3.2 Correct positioning of patient

10.2.3.3 Proper suctioning technique

10.2.3.4 Signs and symptoms of respiratory distress

10.2.3.4 Assessment of patient response to procedure

10.2.3.5 Response to adverse reaction

10.2.3.6 Care and cleaning of equipment

NTS 11.0 MONITORING:

The following should be monitored before, during and following the procedure.

11.1 Breath sounds^{1,3,12,27,59}

11.2 Skin color^{1,6,12,61}

11.3 Breathing pattern and rate^{1,6,12}

11.4 Pulse rate, dysrhythmia, electrocardiogram if available^{1,6,12,14,27,41}

11.5 Color, consistency, and volume of secretions^{1,6}

11.6 Presence of bleeding or evidence of physical trauma^{1,6}

11.7 Subjective response including pain^{1,2,7,41,46}

11.8 Cough¹

11.9 Oxygenation (pulse oximeter)^{1,2,3,6,12,14}

11.10 Intracranial pressure (ICP), if equipment is available¹

11.11 Arterial blood pressure if available⁶

11.12 Laryngospasm⁶

NTS 12.0 FREQUENCY:

Nasotracheal suctioning should be performed by a skilled caregiver when indicated and when other methods to remove secretions from airway have failed.^{1,5,6,8,26,52,54}

NTS 13.0 INFECTION CONTROL:

13.1 CDC Guidelines for Standard Precautions should be adhered to.^{1,12,20,23,60,62}

13.2 All equipment and supplies should be appropriately disposed of or disinfected.^{6,23,62}

Revised by Kim Bennion RRT, Primary Children's Medical Center, Salt Lake City, Utah, and approved by the 2003 CPG Steering Committee

Original Publication: *Respir Care* 1992;37(8):1176-1179.

REFERENCES

1. Burton GG, Hodgkin JE, Ward JJ, editors. *Respiratory care: a guide to clinical practice*, 4th ed. Philadelphia: JB Lippincott; 1997:600-607.
2. Fiorentini A. Potential hazards of tracheobronchial suctioning. *Intensive Crit Care Nurs* 1992;8(4):217-226.
3. Fuchs PL. Streamlining your suctioning techniques. Part I. Nasotracheal suctioning. *Nursing* 1984;14(5):55-61.
4. Demers RR. Management of the airway in the perioperative period. *Respir Care* 1984;29(5):529-536; discussion 536-539.
5. Vender JS, Shapiro BA. Essentials of artificial airway management in critical care. *Acute Care* 1987;13(1-2):97-124.
6. Garvey C. Respiratory therapy function and therapy. In: Nettina SM, editor. *The Lippincott manual of nursing practice*, 7th ed. Philadelphia: JB Lippincott; 2001:221-229.
7. Wanner A, Zigelboim A, Sackner MA. Nasopharyngeal airway: a facilitated access to the trachea: for nasotracheal suction, bedside bronchofiberscopy, and selective bronchography. *Ann Intern Med* 1971;75(4):593-595.
8. Demers RR, Saklad M. Minimizing the harmful effects of mechanical aspiration. *Heart Lung* 1973;2(4):542-545.
9. Griggs A. Tracheostomy: suctioning and humidification. *Nurs Stand* 1998;13(2):49-53.
10. Crow S. Tips for successful respiratory suctioning. *RN* 1986;49(4):31-33.
11. Macmillan C. Nasopharyngeal suction study reveals knowledge deficit. *Nurs Times* 1995;91(50):28-30.
12. Day T, Farnell S, Wilson-Burnett J. Suctioning: a review of current research recommendations. *Intensive Crit Care Nurs* 2002;18(2):79-89.
13. Allen D. Making sense of suctioning. *Nurs Times* 1988;84(10):46-47.
14. Glass CA, Grap MJ. Ten tips for safer suctioning. *Am J Nurs* 1995;95(5):51-53.
15. Plum F, Dunning MF. Techniques for minimizing trauma to the tracheobronchial tree after tracheotomy. *N Engl J Med* 1956;254(5):193-200.
16. James SR, Ashwill JW, Droske SC. *Nursing care of children; principles & practice*, 2nd ed. Philadelphia: WB Saunders; 2002:388.
17. White H. Suctioning: a review. *Paediatr Nurs* 1997;9(4):18-20.
18. Czarnecki ML, Kaucic CL. Infant nasal-pharyngeal suctioning: is it beneficial? *Pediatr Nurs* 1999;25(2):193-196,218.
19. Regan M. Tracheal mucosal injury—the nurse's role. *Nursing (Lond)* 1988;3(29):1064-1066.
20. Day T, Farnell S, Haynes S, Wainwright S, Wilson-Barnett J. Tracheal suctioning: an exploration of nurses' knowledge and competence in acute and high dependency ward areas. *J Adv Nurs* 2002;39(1):35-45.
21. Place B, Fell H. Clearing tracheobronchial secretions using suction. *Nurs Times* 1998;94(47):54-56.
22. Donald KJ, Robertson VJ, Tsebelis K. Setting safe and effective suction pressure: the effect of using a manometer in the suction circuit. *Intensive Care Med* 2000;26(1):15-19.
23. Mathews PJ. 6 steps for safe suctioning. *Nursing* 1994;24(2):18.
24. Boggs RL. Airway management. In: Boggs RL, Woodridge-King M, editors. *AACN procedure manual for critical care*, 3rd ed. Philadelphia: WB Saunders; 1993.
25. Stiller K, Geake T, Taylor J, Grant R, Hall B. Acute lobar atelectasis: a comparison of two chest physiotherapy regimens. *Chest* 1990;98(6):1336-1340.
26. Carroll P. Safe suctioning. *Nursing* 1989;19(9):48-51.
27. Whitaker K. *Comprehensive perinatal and pediatric respiratory care*, 2nd ed. Albany: Delmar Thomson Learning; 1998:211-216.
28. Best Practice. Tracheal suctioning of adults with an artificial airway. *Best Practice* 2000;4(4):1-6.
29. Civeta JM, Taylor RW, Kirby RR. *Critical care*, 3rd ed. Philadelphia: Lippincott-Raven; 1997:1136.
30. Jung RC, Gottlieb LS. Comparison of tracheobronchial suction catheters in humans: visualization by fiberoptic bronchoscopy. *Chest* 1976;69(2):179-181.
31. Landa JF, Kwoka MA, Chapman GA, Brito M, Sackner MA. Effects of suctioning on mucociliary transport. *Chest* 1980;77(2):202-207.
32. Sackner MA, Landa JF, Greenelch N, Robinson MJ. Pathogenesis and prevention of tracheobronchial damage with suction procedures. *Chest* 1973;64(3):284-290.
33. Riegel B, Forshee T. A review and critique of the literature on preoxygenation for endotracheal suctioning. *Heart Lung* 1985;14(5):507-518.
34. Demers RR. Complications of endotracheal suctioning procedures. *Respir Care* 1982;27(4):453-457.

35. Demers RR, Saklad M. Mechanical aspiration: a reappraisal of its hazards. *Respir Care* 1975;20(7):661-666.
36. Touloukian RJ, Beardsley GP, Ablow RC, Effman EL. Traumatic perforation of the pharynx in the newborn. *Pediatrics* 1977;59 Suppl(6 Pt 2):1019-1022.
37. Bogetz MS, Tupper BJ, Vigil AC. Too much of a good thing: uvular trauma caused by overzealous suctioning. *Anesth Analg* 1991;72(1):125-126.
38. Simbruner G, Coradello H, Fodor M, Havelec L, Lubec G, Pollak A. Effect of tracheal suction on oxygenation, circulation, and lung mechanics in newborn infants. *Arch Dis Child* 1981;56(5):326-330.
39. Naigow D, Powaser MM. The effect of different endotracheal suction procedures on arterial blood gases in a controlled experimental model. *Heart Lung* 1977;6(5):808-816.
40. Perlman JM, Volpe JJ. Suctioning in the pre-term infants: effects on cerebral blood flow velocity, intracranial pressure, and arterial blood pressure. *Pediatrics* 1983;72(3):329-334.
41. Brooks D, Anderson CM, Carter MA, Downes LA, Keenan SP, Kelsey CJ, Lacy JB. Clinical practice guidelines for suctioning the airway of the intubated and non-intubated patient. *Can Respir J* 2001;8(3):163-181.
42. Larson RP, Ingalls-Severn KJ, Wright JR, Kiviat NC, Maunder RJ. Diagnosis of *Pneumocystis carinii* pneumonia by respiratory care practitioners: advantages of a nasotracheal suctioning method over sputum induction. *Respir Care* 1989;34(4):249-253.
43. Fox WW, Schwartz JG, Shaffer TH. Pulmonary physiotherapy in neonates: physiologic changes and respiratory management. *J Pediatr* 1978;92(6):977-981.
44. Wilkins RL, Stoller JK, Scanlan CL, editors. Egan's fundamentals of respiratory care, 8th ed. St Louis: CV Mosby; 2003.
45. Cordero L Jr, Hon EH. Neonatal bradycardia following nasopharyngeal stimulation. *J Pediatr* 1971;78(3):441-447.
46. Brucia J, Rudy E. The effect of suction catheter insertion and tracheal stimulation in adults with severe brain injury. *Heart Lung* 1996;25(4):295-303.
47. Fisher DM, Frewen T, Swedlow DB. Increase in intracranial pressure during suctioning—stimulation vs rise in P_{aCO_2} . *Anesthesiology* 1982;57(5):416-417.
48. Parsons LC, Shogan JS. The effects of the endotracheal tube suctioning/manual hyperinflation procedure on patients with severe closed head injuries. *Heart Lung* 1984;13(4):372-380.
49. Ersson U, Carlson H, Mellström A, Pontén U, Hedstrand U, Jakobsson S. Observations on intracranial dynamics during respiratory physiotherapy in unconscious neurosurgical patients. *Acta Anaesthesiol Scand* 1990;34(2):99-103.
50. Evans JC. Reducing the hypoxemia, bradycardia, and apnea associated with suctioning in low birthweight infants. *J Perinatol* 1992;12(2):137-142.
51. Petersen GM, Pierson DJ, Hunter PM. Arterial oxygen saturation during nasotracheal suctioning. *Chest* 1979;76(3):283-287.
52. Hoffman LA, Maszkiewicz RC. Airway management for the critically ill patient. *Am J Nurs* 1987;87(1):39-53.
53. Day T. Tracheal suctioning: when, why and how. *Nursing Times* 2000;96(20):13-15.
54. Petty TL. Intensive and rehabilitative respiratory care, 3rd ed. Philadelphia: Lea & Febiger; 1982:30-33,154.
55. Curran JF, Stanek KS, Kacmarek RM. Portable airway-suction systems: a comparison of performance. *Respir Care* 1991;36(4):259-266.
56. Das PK, Thomas WJ. Complication of pharyngeal suction. *Anaesth Intensive Care* 1980;8(3):375-376.
57. Langrehr EA, Woodburn SC, Guthrie MP. Oxygen insufflation during endotracheal suctioning. *Heart Lung* 1981;10(6):1028-1036.
58. Jacqueline G. To reduce hazards of tracheal suctioning. *Am J Nurs* 1971;71(12):2362-2364.
59. Occupational exposure to blood borne pathogens. Final Rule. Federal Register 1991;56:64175-64182.
60. Bolyard EA, Tablan OC, Williams WW, Pearson ML, Shapiro CN, Deitchmann SD. Guideline for infection control in healthcare personnel, 1998. Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 1998;19(6):407-463. Erratum in: *Infect Control Hosp Epidemiol* 1998;19(7):493.
61. Update: universal precautions for prevention of transmission of human immunodeficiency virus, hepatitis B virus, and other bloodborne pathogens in health-care settings. *MMWR Morb Mortal Wkly Rep* 1988;37(24):377-382, 387-388.
62. Creamer E, Smyth EG. Suction apparatus and the suctioning procedure: reducing the infection risks. *J Hosp Infect* 1996;34(1):1-9.

Interested persons may photocopy these Guidelines for noncommercial purposes of scientific or educational advancement. Please credit AARC and RESPIRATORY CARE Journal.

All of the AARC CPGs may be downloaded at no charge from
<http://www.rcjournal.com/online/>